SE 461 Computer Vision

Nazar Khan

PUCIT


Lecture 3

Topics to be covered

- Motivation for Studying Colour
- Physical Background
- Biological Background
- Technical Colour Spaces


Motivation

- Colour science or colour vision
 - highly interdisciplinary area
 - influences from physics, biology, psychology, physiology, electrical engineering, computer science, and mathematics
- Understanding the mechanisms of colour perception is crucial for
 - calibrating scanners, colour monitors, colour printers, and digital cameras
- Many brilliant minds have done research on colour
 - Newton, Goethe, Graßmann, Maxwell, Helmholtz, Schroedinger.
 - Not always successful




The wave length λ is the distance between two consecutive points with the same phase. Source: http://en.wikipedia.org/wiki/Wavelength

- Electromagnetic waves propagate with the speed of light
 - -c = 299, 792, 458 m/s in vacuum.
 - If f denotes their frequency, wave length λ is defined via $c = \lambda f$.
- Visible light consists of electromagnetic waves with wave lengths λ between 380 and 750 nm (nanometers)
 - $-1 nm = 10^{-9} m$
- Light is emitted from atoms when outer electrons change their orbits and loose energy.




Visible light constitutes only a small part of the entire electromagnetic spectrum. Author: N. Khan (2005).




- Visible range of the electromagnetic spectrum is the main range in which the Sun's rays can pass the atmosphere
- The visible range is not the same for all animals
 - Insects can perceive ultraviolet light very well
 - Humans whose eye lenses have been removed by a surgeon also perceive ultraviolet light down to 300 nm
- The visible spectrum contains seven colours
 - red, orange, yellow, green, blue, indigo, violet
 - with decreasing wavelength or increasing frequency

- Colours that we usually perceive are a mixture of numerous frequencies.
- Pure spectral colours consisting of a single frequency are rare
 - They exist e.g. in rainbows, in light rays that pass through a prism, and in lasers.



Source: http://www.theinquirer.net/IMG/847/948 47/light-split-into-spectrum-by-prism.jpg


- However, biological and technical systems do not exploit all frequencies of a perceived colour
 - They use a more compact representation.
- To understand this, let us consider the biology of colour perception.



The Eye
http://www.soundandvision.com/images/
archivesart/204eye.3.jpg

The Retina
http://www.pennmedicine.org/health-info/
body-guide/reftext/images/retina-section.jpg

- The human retina consists of two classes of photo receptors:
 - 120 million rods
 distributed over the
 whole retina. Multiple
 rods are connected to a
 single nerve.
 - 6 million cones, mainly in the centre of the retina.
- Each cone is connected to its own nerve end.




*ADAM

The Retina

http://www.pennmedicine.org/health info/
body guide/reftext/images/retina section.jpg

- For night vision, rods dominate.
 - Very sensitive, but allows only poor sharpness and no colour perception
- Day vision is dominated by cones.
 - Low sensitivity, but very good perception of sharpness and colours.
- Humans can distinguish about 40 different grayscales, but 2 million colours.
- Our focus day vision.

- Humans have three different types of cones (L, M, S) being sensitive to different wavelength ranges:
 - Long → red
 - Medium → green
 - Medium → blue.
- The sensitivity characteristics are almost identical for all humans.



Normalised spectral sensitivity of the three cone types in the human eye.

Author: M. Mainberger (2008).

- Visual impression of a colour depends on its effect on the 3 types of cones.
 - Specifically: The colour produces 3 response values (one for each kind of cone)
 - These are called tri-stimulus values
- Mathematically speaking


The tristimulus value Q_i of a cone of type $i=1,\,2,\,3$ for L-, M-, and S-cones, respectively, is given by

$$Q_i = \int C(\lambda) R_i(\lambda) d\lambda$$

where

- ullet λ denotes the wave length,
- $C(\lambda)$ is the incoming stimulus (product of illuminant and reflectance),
- $R_i(\lambda)$ is the spectral sensitivity of cone i.

$$Q_S = \int \text{Incoming_light}(\lambda) \cdot \text{Blue_Curve}(\lambda) d\lambda$$





- Colours are no objective physical reality.
- Colour perception is a biologically useful interpretation of the physically existing wave length spectrum.

The RGB Colour Space

- Additive Model
- Primary colour basis
 - red (R = (1, 0, 0))
 - green (G = (0, 1, 0))
 - blue (B = (0, 0, 1))
- All colors are generated as linear combinations of these 3 primary colours.
 - This yields the so-called RGB Cube

- Every colour has a complementary colour such that adding them gives white.
- The complementary colour is at the opposite side of the cube.






Colours can be blended by linear combinations of the primary colors in the RGB space. Computer screens use this kind of **additive** colour blending.

The CMY Color Space

- Used by printers and copiers that place colour pigments on <u>white</u> paper.
- In contrast to the additive RGB model, the CMY model is subtractive.
- Colour impressions are determined by the colour that is subtracted (absorbed into the white paper) from white light.
- Complementary to RGB
 - Basis: cyan (C=(1,0,0)), magenta (M=(0,1,0)), and yellow (Y=(0,0,1))
- In the RGB model, an equal combination of the primary colours R, G, B gives white.
- In the CMY model, an equal combination of the primary colours C, M, Y creates ... ?
 - black.



$$\left(\begin{array}{c} C \\ M \\ Y \end{array}\right) = \left(\begin{array}{c} 1 \\ 1 \\ 1 \end{array}\right) - \left(\begin{array}{c} R \\ G \\ B \end{array}\right)$$


Colours can be blended by linear combinations of the primary colors in the CMY space. Printers use this kind of **subtractive** colour blending.

The CMYK Colour Space

- Extension of CMY to include black (K) as fourth colour
- Useful for colour printers that print large amounts of text in black
- Conversion formula for CMY to CMYK

$$K \leftrightarrow \min(C, M, Y)$$
 $C \leftrightarrow C - K$
 $M \leftrightarrow M - K$
 $Y \leftrightarrow Y - K$

The HSV Colour Space



The HSV colour space uses a cylindrical polar coordinate system with hue, saturation and value as coordinates. Source: http://de.wikipedia.org/wiki/HSV-Farbraum.

The HSV Colour Space


- Based on a cylindrical polar coordinate system with hue, saturation and value as coordinates
- Hue (H)
 - polar angle in the horizontal plane
 - red: 0; green: 120; blue: 240
- saturation (S)
 - radius in the horizontal plane
 - gives distance of the colour to the next grey tone
- value (V)
 - vertical axis in the coordinate system
 - defines how dark or bright a colour is

HSV can offer advantages when illumination changes appear

- Typical changes
 - global multiplicative changes: global illumination changes
 - local multiplicative changes: shadow, shading
 - local additive changes: specular highlights
- The H and S channels are photometric invariants:
 - H is invariant under all these three changes.
 - S is invariant under global and local multiplicative changes.
 - V is not invariant under any of these changes.

The YCbCr Colour Space

- Used in video and digital photography systems
 - digital PAL and NTSC television,
 - JPEG and MPEG compression
- separates the colour image into
 - a luma component Y that is basically a greyscale version of the colour image
 - a chroma component Cb that measures deviation from grey in blue—yellow direction
 - a chroma component Cr that measures deviation from grey in red-cyan direction



Top left: Original image, 512×512 pixels. Top right: The luma channel Y contains the luminance information with lots of details. Bottom left: Blue-yellow chroma channel Cb. Bottom right: Red-cyan chroma channel Cr. Author: M. Mainberger (2009).

Color Spaces and Image Compression

- The luma channel Y should be stored in high resolution, while the chroma channels Cb and Cr can be subsampled without significant visual deterioration.
- This is exploited in the JPEG compression standard.

Summary

- Visible light consists of electromagnetic waves in the range from 380 to 750 nanometers.
- Illuminated real objects emit a mixture of different wave lengths.
- Human colour perception is based on three types of cones that are sensitive in different frequency bands.
- An additive blending of the three primary colours red, green and blue is used for creating other colours.
- This is realised in the RGB colour space which is a very frequently used additive colour space for digital cameras and monitors.

Summary

- The CMY and CMYK spaces are subtractive colour spaces for printers and copiers.
- The HSV colour space represents colours in a cylindrical polar coordinate system.
 - This can be useful when illumination changes appear.
- The YCbCr representation separates luma from chroma information.
 - One can subsample the chroma channels without severe visual degradation.
 - Useful for image compression.